

GROUP	1	HERBICIDE
-------	----------	-----------

FOOTHILLS™ A Herbicide

EMULSIFIABLE CONCENTRATE

AGRICULTURAL

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND PEACE RIVER, OKANAGAN AND CRESTON FLATS REGIONS OF BRITISH COLUMBIA ONLY.

A post emergence herbicide for control of wild oats, volunteer (tame) oats, green foxtail, yellow foxtail, barnyard grass, Persian darnel and volunteer canary seed in spring wheat and Durum wheat.

GUARANTEE:

Clodinafop-propargyl..... 240 g/L

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING.
KEEP OUT OF REACH OF CHILDREN.**

CAUTION



POISON

WARNING: EYE & SKIN IRRITANT

Warning, contains the allergen epoxidized soybean oil

REGISTRATION NO.: 29279
PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1.84L, 3.68L, 4.7L, 14L, 15L, 55L, 450L, 1100L, Bulk

Syngenta Crop Protection Canada, Inc.

140 Research Lane, Research Park
Guelph, Ontario
N1G 4Z3
Telephone: 1-877-964-3682

LABEL

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

TOXICOLOGICAL INFORMATION

This product contains materials that may cause severe pneumonitis if aspirated. Treat symptomatically for ingestion and/or skin and eye contact. This product contains PETROLEUM DISTILLATES. DO NOT INDUCE VOMITING.

PRECAUTIONS

- CAUTION – POISON
- WARNING – EYE AND SKIN IRRITANT
- KEEP OUT OF THE REACH OF CHILDREN.
- Harmful if swallowed.
- May irritate eyes. Do not wear contact lenses when using.
- DO NOT get in eyes or on skin. Avoid contact with clothing. Wear coveralls or long sleeved shirt and long pants, chemical resistant gloves, and goggles when mixing, loading or during equipment clean up or repair.
- Wash gloves thoroughly with soap and water before removing them during any operation.
- Wash hands thoroughly with soap and water after using this product and before eating, drinking or smoking.

- Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before reuse. Wash thoroughly with soap and water after handling. Handle and apply only as recommended on this label.
- Do not eat, drink or smoke while mixing, loading or during application.
- Do not contaminate streams, rivers, lakes, ponds, wetlands, sloughs, irrigation water, vegetated ditch banks, and water used by livestock or for domestic purposes.
- This product contains a petroleum distillate, which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal of waste or cleaning of equipment.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in U.S., visit CropLife Canada's website at www.croplife.ca.

STORAGE

Store the product in closed original container in a well-ventilated room. Keep out of reach of children, unauthorized persons and animals. Store separate from food, feed, and fertilizer.

DISPOSAL OF UNUSED, UNWANTED PRODUCT

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

CONTAINER DISPOSAL:

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

GROUP 1 HERBICIDE

FOOTHILLS™ A Herbicide

EMULSIFIABLE CONCENTRATE

AGRICULTURAL

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND PEACE RIVER, OKANAGAN AND CRESTON FLATS REGIONS OF BRITISH COLUMBIA ONLY.

A post emergence herbicide for control of wild oats, volunteer (tame) oats, green foxtail, yellow foxtail, barnyard grass, Persian darnel and volunteer canary seed in spring wheat and Durum wheat.

GUARANTEE:

Clodinafop-propargyl..... 240 g/L

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING.
KEEP OUT OF REACH OF CHILDREN.**

CAUTION



POISON

WARNING: EYE & SKIN IRRITANT

Warning, contains the allergen epoxidized soybean oil

REGISTRATION NO.: 29279
PEST CONTROL PRODUCTS ACT

Syngenta Crop Protection Canada, Inc.

140 Research Lane, Research Park
Guelph, Ontario
N1G 4Z3
Telephone: 1-877-964-3682

PAMPHLET

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

TOXICOLOGICAL INFORMATION

This product contains materials that may cause severe pneumonitis if aspirated. Treat symptomatically for ingestion and/or skin and eye contact. This product contains PETROLEUM DISTILLATES. DO NOT INDUCE VOMITING.

PRECAUTIONS

- CAUTION – POISON
- WARNING – EYE AND SKIN IRRITANT
- KEEP OUT OF THE REACH OF CHILDREN.
- Harmful if swallowed.
- May irritate eyes. Do not wear contact lenses when using.
- DO NOT get in eyes or on skin. Avoid contact with clothing. Wear coveralls or long sleeved shirt and long pants, chemical resistant gloves, and goggles when mixing, loading or during equipment clean up or repair.
- Wash gloves thoroughly with soap and water before removing them during any operation.
- Wash hands thoroughly with soap and water after using this product and before eating, drinking or smoking.

- Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before reuse. Wash thoroughly with soap and water after handling. Handle and apply only as recommended on this label.
- Do not eat, drink or smoke while mixing, loading or during application.
- Do not contaminate streams, rivers, lakes, ponds, wetlands, sloughs, irrigation water, vegetated ditch banks, and water used by livestock or for domestic purposes.
- This product contains a petroleum distillate, which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal of waste or cleaning of equipment.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in U.S., visit CropLife Canada's website at www.croplife.ca.

STORAGE

Store the product in closed original container in a well-ventilated room. Keep out of reach of children, unauthorized persons and animals. Store separate from food, feed, and fertilizer.

DISPOSAL OF UNUSED, UNWANTED PRODUCT

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

CONTAINER DISPOSAL:

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

PRODUCT INFORMATION

FOOTHILLS A Herbicide is a systemic, post-emergence herbicide for the selective control of wild oats, volunteer (tame) oats, green foxtail, yellow foxtail, barnyard grass, Persian darnel and volunteer canary seed in all types of spring wheat and Durum wheat. Do not use FOOTHILLS A Herbicide on barley, as crop injury will occur. Do not apply this product using aerial application equipment except under conditions specified on this label.

FOOTHILLS A Herbicide is absorbed by the leaves and is rapidly translocated to the growing points of leaves and stems. Thorough coverage of the plants is essential for consistent control. Actively growing susceptible grasses stop growing within 48 hours of treatment. Depending on species, growing conditions and crop competition, leaves and growing points turn yellow within one to three weeks after application. Further colour changes and loss of vigour will be observed, followed by a browning and complete control three to five weeks after application.

Although FOOTHILLS A Herbicide does not control broadleaf weeds, FOOTHILLS A Herbicide can be tank-mixed with a wide range of broadleaf herbicides to provide broad spectrum weed control in wheat. It can also be tank mixed with Decis[†] Flowable Insecticide and MATADOR[®] 120EC Insecticide to control grassy weeds and insects and with TILT[®] 250E Fungicide to control grassy weeds and cereal leaf diseases. See section entitled "TANK MIXES of FOOTHILLS A HERBICIDE WITH BROADLEAF WEED HERBICIDES, INSECTICIDES AND FUNGICIDES" and refer to the appropriate Tank Mix section for ground and aerial application.

DIRECTIONS FOR USE

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted, fine textured, or low in organic matter such as clay.)

Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip (buffer zone) between the treated area and the edge of the water body.

CROPS: Spring Wheat and Durum Wheat

FOOTHILLS A Herbicide can be used on all varieties of spring wheat and Durum wheat. Observe minimum interval to harvest of 60 days after treatment.

Do not apply on barley or any crop other than Spring Wheat or Durum wheat, as crop damage will result. Do not allow spray to drift to adjacent fields seeded to crops other than spring wheat or Durum wheat.

Do not treat wheat underseeded to forages. Observe a minimum of three (3) days before grazing livestock on crops treated with FOOTHILLS A Herbicide.

For use directions specific to application by air, please refer to sections on aerial application. For aerial application precautions, please refer to the precautions section.

WEEDS CONTROLLED: Wild Oats, Volunteer (Tame) Oats, Green Foxtail, Yellow Foxtail, Barnyard grass, Persian Darnel and Volunteer Canary Seed

TIMING OF APPLICATION:

TIMING	GROWTH STAGE	ADDITIONAL REMARKS
WILD OATS	1 to 6 leaf stage on main stem	Prior to emergence of 4th tiller
VOLUNTEER (TAME) OATS	3 to 6 leaf stage on main stem	Prior to emergence of 4th tiller
GREEN FOXTAIL AND YELLOW FOXTAIL (wild millet, pigeon grass)	1 to 5 leaf stage on main stem	For optimum control apply prior to emergence of the 3 rd tiller and while foxtail is actively growing
BARNYARD GRASS	1 to 5 leaf stage on main stem	For optimum control apply before tillering, and while barnyard grass is actively growing
PERSIAN DARNEL	1 to 5 leaf stage on main stem	For optimum control apply before tillering, and while Persian darnel is actively growing
VOLUNTEER CANARY SEED	1 to 6 leaf stage on main stem	Prior to emergence of 4th tiller
SPRING WHEAT AND DURUM WHEAT	Prior to emergence of 4th tiller	When tank-mixing with a broadleaf herbicide, insecticide or a fungicide, always refer to the label of the broadleaf partner, insecticide partner or fungicide partner prior to use.

- For optimum results, apply FOOTHILLS A Herbicide to actively growing weeds. An early application will maximize crop yields by reducing weed competition. Weeds emerging after application of FOOTHILLS A Herbicide will not be controlled.
- Weed control following application of FOOTHILLS A Herbicide alone, or in combination with broadleaf weed herbicides, can be reduced or delayed under stress conditions such as drought, heat, insufficient fertility, flooding or prolonged cool temperatures. Grass escapes or re-tillering may occur if application is made during prolonged stress conditions. Optimum weed control will be obtained if application of FOOTHILLS A Herbicide is delayed until the stress conditions have ended and weeds are once again actively growing.
- FOOTHILLS A Herbicide alone can be used 30 minutes before rainfall.
- Do not apply to crop that is stressed by conditions such as frost, low fertility, drought, flooding, disease or insect damage as crop injury may result.

RATE OF USE FOR GROUND APPLICATION:

Apply the recommended rate of Foothills A Herbicide and the recommended rate of Foothills™ B Adjuvant in a minimum of 50 to 100 L of water per hectare.

<p>To control:</p> <p>WILD OATS VOLUNTEER (TAME) OATS GREEN FOXTAIL YELLOW FOXTAIL BARNYARD GRASS VOLUNTEER CANARY SEED</p>	<p>To control:</p> <p>WILD OATS VOLUNTEER (TAME) OATS GREEN FOXTAIL YELLOW FOXTAIL BARNYARD GRASS VOLUNTEER CANARY SEED PERSIAN DARNEL</p>
Rates for use with 50 L/ha water	
<p>Apply:</p> <p>230 mL/ha of Foothills A Herbicide + 400 mL/ha of Foothills B Adjuvant with 50 L/ha of water (0.8% volume/volume)</p>	<p>Apply:</p> <p>290 mL/ha of Foothills A Herbicide + 500 mL/ha of Foothills B Adjuvant with 50 L/ha of water (1.0% volume/volume)</p>
<p>NOTE:</p> <p>One 1.84/3.68/4.0/4.7/8.0 L jug of Foothills A Herbicide + 3.2/6.4/6.96/8.16/14.0 L Foothills B Adjuvant treats 8 ha (20 acres)/16 ha (40 acres)/17.4 ha (43 acres)/20.4 ha (50.4 acres)/35 ha (86.4 acres) with 400/800/870/1020/1750 L of water.</p>	<p>NOTE:</p> <p>One 1.84/3.68/4.0/4.7/8.0 L jug of Foothills A Herbicide + 3.2/6.4/6.96/8.16/14.0 L Foothills B Adjuvant treats 6.4 ha (15.8 acres)/12.8 ha (31.4 acres)/13.9 ha (34.3 acres)/16.3 ha (40.2 acres)/28 ha (69 acres) with 320/640/695/815/1400 L of water.</p>
Rates for use with 100 L/ha water	
<p>Apply:</p> <p>230 mL/ha of Foothills A Herbicide + 800 mL/ha of Foothills B Adjuvant with 100 L/ha of water (0.8% volume/volume)</p>	<p>Apply:</p> <p>290 mL/ha of Foothills A Herbicide + 1 L/ha of Foothills B Adjuvant with 100 L/ha of water (1.0% volume/volume)</p>
<p>NOTE:</p> <p>One 1.84/3.68/4.0/4.7/8.0 L jug of Foothills A Herbicide + 6.4/12.8/13.9/16.3/28.0 L Foothills B Adjuvant treats 8 ha (20 acres)/16 ha (40 acres)/17.4 ha (43 acres)/20.4 ha (50.4 acres)/35 ha (86.4 acres) with 800/1600/1740/2040/3500 L of water.</p>	<p>NOTE:</p> <p>One 1.84/3.68/4.0/4.7/8.0 L jug of Foothills A Herbicide + 6.4/12.8/13.9/16.3/28.0 L Foothills B Adjuvant treats 6.4 ha (15.8 acres)/12.8 ha (31.4 acres)/13.9 ha (34.3 acres)/16.3 ha (40.2 acres)/28 ha (69 acres) with 640/1280/1390/1630/2800 L of water.</p>

NOTE: Always use Foothills B Adjuvant with Foothills A Herbicide.

RATE OF USE FOR AERIAL APPLICATION:

Apply the recommended rate of Foothills A Herbicide and the recommended rate of Foothills B Adjuvant in a minimum of 30 L of water per hectare according to the following table:

To Control: WILD OATS	To Control: WILD OATS GREEN FOXTAIL YELLOW FOXTAIL PERSIAN DARNEL
Apply: 230 mL/ha of Foothills A Herbicide + 240 mL/ha of Foothills B Adjuvant with 30 L/ha of water (0.8% volume/volume)	Apply: 290 mL/ha of Foothills A Herbicide + 300 mL/ha of Foothills B Adjuvant with 30 L/ha of water (1.0% volume/volume)
NOTE: <i>One 1.84/3.68/4.0/4.7/8.0 L jug of Foothills A Herbicide + 1.92/3.84/4.18/4.9/8.4 L Foothills B Adjuvant treats 8 ha (20 acres)/16 ha (40 acres)/17.4 ha (43 acres)/20.4 ha (50.4 acres)/35 ha (86.4 acres) with 240/480/522/612/1050 L of water.</i>	NOTE: <i>One 1.84/3.68/4.0/4.7/8.0 L jug of Foothills A Herbicide + 1.92/3.84/4.17/4.9/8.4 L Foothills B Adjuvant treats 6.4 ha (15.8 acres)/12.8 ha (31.4 acres)/13.9 ha (34.3 acres)/16.3 ha (40.2 acres)/28 ha (69 acres) with 192/384/417/490/840 L of water.</i>

NOTE: Always use Foothills B Adjuvant with Foothills A Herbicide.

TANK MIXES of Foothills A Herbicide with Broadleaf Weed Herbicides, Insecticides and Fungicides – Ground Application:

CROP: SPRING and DURUM WHEAT

TANK MIXES WITH Broadleaf Weed Herbicides, Insecticides and Fungicides in Spring Wheat and Durum Wheat:

For broad spectrum control of wild oats, green foxtail, barnyard grass, Persian Darnel and broadleaf weeds, Foothills A Herbicide Tank Mix can be tank-mixed with broadleaf herbicides as described in the following tables. Foothills A Herbicide Tank Mix can also be tank-mixed with Decis[†] Flowable Insecticide and MATADOR[®] 120EC Insecticide for control of insects listed on the Decis Flowable Insecticide and MATADOR 120EC Insecticide label and with TILT 250E Fungicide for control of diseases listed on the TILT[®] 250E Fungicide label, in addition to grassy weeds. Consult the label of the tank-mix partner for a list of broadleaf weeds, insects or diseases controlled, rates, timing, recropping restrictions, grazing interval restrictions, recommendations for specific weeds/insects/diseases, directions for use and precautions and follow the more restrictive label. When tank-mixing always add the broadleaf herbicide(s), insecticide or fungicide to the spray tank first; followed by Foothills A Herbicide, with Foothills B Adjuvant added last. For the appropriate rate of Foothills A Herbicide with Foothills B Adjuvant, refer to the 'Rate of Use' section of the label.

Tank mixes for broad spectrum control of wild oats, green foxtail, broadleaf weeds, insects and diseases:

Tank-Mix Partner	Product Rates	Crop Stage¹
TARGET [®]	1.0 to 1.5 L/ha	2 to 5 leaf
Dyvel [†]	1.25 L/ha	2 to 5 leaf
Refine Extra ^{†2}	20 g/ha	2 leaf to flag leaf
Buctril [†] M	1.0 L/ha	2 leaf to flag leaf
Estaprop [†]	1.75 L/ha	4 leaf to early flag leaf (shot blade)
Turboprop [†] 600	1.75 L/ha	4 leaf to early flag leaf (shot blade)
Dichlorprop [™] -D	1.75 L/ha	4 leaf to early flag leaf (shot blade)
Unity ^{†2}	1 carton/16 ha	2 leaf to flag leaf
Lontrel [†] 360EC	0.42 to 0.84 L/ha	3 leaf to flag leaf
Lontrel 360EC tank mixed with MCPA Ester (assume 500 series)	280 - 420 mL/ha tank mixed with 1.1 L/ha	3 leaf to flag leaf
Lontrel 360EC tank mixed with Refine Extra 75DF	0.21 L/ha tank mixed with 20 g/ha	3 leaf to flag leaf
Curtail [†] M	1.5 to 2.0 L/ha	3 leaf to just before flag leaf
Prestige [†] Herbicide Tank Mix ⁹	600 - 800 mL/ha of Prestige A + 1.5 - 2.0 L/ha of Prestige B	3 leaf to flag leaf
Thumper [®] EC	1.0 L/ha	4 leaf to flag leaf
2,4-D Amine (assume 500 series) ³	840 mL to 1.1 L/ha	3 leaf to flag leaf
MCPA Amine (assume 500 series) ³	840 mL to 1.1 L/ha	3 leaf to flag leaf
MCPA Ester (assume 500 series)	840 mL to 1.1 L/ha	3 leaf to flag leaf
MCPA 300SL Sodium salt	1.2 L to 2.7 L/ha ⁵	3 leaf to flag leaf
Pardner [†]	1.0 L/ha	2 leaf to flag leaf
Ally ^{†2}	7.5 g/ha	2 leaf to flag leaf
Attain [†] Herbicide Tank Mix ⁹	450 - 600 mL/ha of Attain A + 0.75 - 1.0 L/ha of Attain B	4 leaf to flag leaf
Decis Flowable Insecticide	80-120 mL/ha	Prior to emergence of 4 th tiller
TILT [®] 250E Fungicide	500 mL/ha	Prior to emergence of 4 th tiller
Mecoprop ⁵	5.5 - 7.0 L/ha	3 leaf to just before flag leaf
MATADOR [®] 120EC Insecticide	63 - 83 mL/ha	Prior to emergence of 4 th tiller
Mextrol [†] 400M	1.4 L/ha	2 leaf to flag leaf
MEXTROL 450 Liquid Herbicide	1.25 L/ha	2 leaf to flag leaf
Trophy Herbicide Tank-Mix ⁴	0.6 L/ha of Trophy A + 1.12 L/ha Trophy B	3 leaf to flag leaf
Florasulam [†] SC Herbicide Tank Mix	100 mL/ha of Florasulam SC	2 to 4 leaf
Florasulam SC Herbicide + MCPA LV 500 Tank Mix ⁶	100 mL/ha of Florasulam SC + 0.84 L/ha of MCPA LV 500	2 to 6 leaf
Frontline [†] Herbicide Tank Mix ^{6,7}	100 mL/ha of Frontline A + 700 mL/ha of Frontline B	2 to 6 leaf
Refine Extra ^{†2} tank-mixed with Accord [†] Water-Soluble Bag Herbicide ⁸	20 g/ha of Refine Extra + 67 g/ha of Accord W.S.B.	2 to 5 leaf
Refine SG [†] Herbicide ²	30 g/ha	2 leaf to flag leaf
Refine SG Toss-N-Go Herbicide ^{†2}	30 g/ha	2 leaf to flag leaf
DYCLEER Agricultural Herbicide + Starane Herbicide ¹⁰	(0.11 + 0.39L/ha) to (0.16 + 0.58 L/ha)	2-5 leaf
¹ Always consult the label of the broadleaf herbicide, insecticide or fungicide prior to use.		
² Addition of surfactants other than FOOTHILLS B is not required.		
³ A reduction in control of green foxtail and wild oats may be observed when FOOTHILLS A Herbicide is tank mixed with 2,4-D Amine and MCPA Amine.		
⁴ Rates above 2.0 L/ha may cause crop injury		
⁵ Tank mix with Mecoprop provides suppression of Canada Thistle (top growth control)		

⁶ Use with the high rate of Foothills A Herbicide for wild oat and green foxtail control.
⁷ Use additional 140mL of Frontline B for optimum control of: Redroot Pigweed, Hempnettle, Dandelion, Stork's Bill, and Canada Thistle.
⁸ When tank-mixing with Accord Water-Soluble Bag Herbicide, do not graze treated crops or cut for hay within 77 days of application.
⁹ Refer to broadleaf tank-mix label for list of weeds controlled at low and high use rates.
¹⁰ When tank-mixing with DYCLEER +Starane use Foothills B Adjuvant.

For the control of barnyard grass and broadleaf weeds, in addition to wild oats and green foxtail, use one of the following tank mixes:

Tank Mix Partner	Product Rates	Crop Stage ¹
Target	1.0 to 1.5 L/ha	2 to 5 leaf
Buctril M	1.0 L/ha	2 leaf to flag leaf
Estoprop	1.75 L/ha	4 leaf to early flag leaf (shot blade)

¹ Always consult the label of the broadleaf herbicide prior to use.

For control of Persian Darnel* and broadleaf weeds, in addition to wild oats, green foxtail and barnyard grass, use the following tank mix:

Tank Mix Partner	Product Rates	Crop Stage ¹
Buctril M	1.0 L/ha	2 leaf to flag leaf

¹ Always consult the label of the broadleaf herbicide prior to use.

* Use the higher rate of Foothills A Herbicide Tank Mix to control Persian Darnel. Refer to the "Rate of Use" section of the label.

Temporary crop injury may occur with tank-mixes under extreme weather conditions or when the crop is suffering from stress due to inadequate or abnormally high moisture level or extreme temperatures.

Do not tank-mix with any chemical additives, pesticides, or fertilizers that are not recommended on this label.

TANK MIXES OF Foothills A Herbicide WITH Broadleaf Weed Herbicides - Aerial Application:

CROP: SPRING WHEAT and DURUM WHEAT

For broad spectrum control of wild oats, green foxtail and broadleaf weeds, Foothills A Herbicide can be tank-mixed with Buctril M. Consult the label of the tank-mix partner for a list of broadleaf weeds controlled, rates, timing, recropping restrictions, grazing interval restrictions, recommendations for specific weeds, directions for use and precautions and follow the more restrictive label. Use in a minimum of 30L of water per hectare.

Tank-mixes of Foothills A Herbicide with broadleaf weed herbicides - Aerial application:
When tank-mixing always add the broadleaf herbicide (Buctril M) to the spray tank first; followed by Foothills A Herbicide, with Foothills B Adjuvant added last. For the appropriate rate of Foothills A Herbicide with Foothills B refer to the 'Rate of Use For Aerial Application' section of the label.

Tank-Mix Partner	Product Rates	Crop Stage ¹
Buctril M	1.0 L/ha	2 leaf to flag leaf

¹ Always consult the label of the broadleaf partner prior to use.

Temporary crop injury may occur with tank-mixes under extreme weather conditions or when the crop is suffering from stress due to inadequate or abnormally high moisture level or extreme temperatures.

Do not tank-mix with any chemical additives, pesticides, or fertilizers that are not recommended on this label.

MIXING INSTRUCTIONS - GROUND APPLICATION:

1. Clean spray tank and half fill with clean water. Start agitation or bypass system
2. If a broadleaf herbicide, insecticide or fungicide is to be used, add the product FIRST prior to adding FOOTHILLS A Herbicide and agitate for 2-3 minutes.
3. Add correct amount of FOOTHILLS A Herbicide.
4. Agitate for 2-3 minutes.
5. Add correct amount of FOOTHILLS B Adjuvant.
6. Agitate for 1-2 minutes before adding remainder of water and then maintain constant agitation.
7. After any break in spraying operations, agitate thoroughly before spraying again.
8. **Use the spray suspension as soon as it is prepared.**
9. If an oil film starts to build up in the tank, drain tank and then clean with a detergent.

SPRAYING INSTRUCTIONS - GROUND APPLICATION:

1. FOOTHILLS A can be applied by ground or air. For aerial application instructions, refer to the section entitled "SPRAYING INSTRUCTIONS - AERIAL APPLICATION" which follows this section.
2. Water Volume: 50 to 100 litres per hectare when applied alone and a minimum of 100L/ha when tank-mixed with broadleaf herbicides
3. Spray Nozzles: 80° or 110° flat fan stainless steel nozzles are recommended for optimal spray coverage. Application of the spray mixture at a 45° angle in the direction of travel will result in improved spray coverage. Use 50 mesh nozzle screens. Do not use flood type nozzles, controlled droplet application equipment, spray foils or hollow cone nozzles.
4. Pressure: 275-310 kPa.
5. Apply uniformly at 6-8 km/hr and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from an over application.
6. Do not contaminate streams, rivers, lakes, ponds, wetlands, sloughs, irrigation water, vegetated ditch banks, and water used by livestock or for domestic purposes. Do not apply within 15 meters of a wetland area or any of the aforementioned bodies of water.

AERIAL APPLICATION

Generic Aerial Application Label Instructions - Directions for Use

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on

this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-87-SYNGENTA (1-877-964-3682) or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

MIXING INSTRUCTIONS - AERIAL APPLICATION:

1. Fill the mixing tank 1/2 full with clean water. Start gentle agitation.
2. If a broadleaf herbicide is to be used, add the product **FIRST** prior to adding Foothills A Herbicide and agitate for 2-3 minutes.
3. Add correct amount of Foothills A Herbicide.
4. Agitate for 2-3 minutes.

5. Add correct amount of FOOTHILLS B Adjuvant.
6. Agitate for 1-2 minutes before adding remainder of water and then maintain constant agitation.
7. Fill aircraft spray tank and maintain gentle agitation while spraying.
8. After any break in spraying operations, agitate thoroughly before spraying again. Do not let contents stand without agitation.
9. **Use the spray suspension as soon as it is prepared.**
10. If an oil film starts to build up in the tank, drain tank and then clean with a detergent.

SPRAYING INSTRUCTIONS - AERIAL APPLICATION:

1. FOOTHILLS A Herbicide can be applied by ground or by air. For ground application instructions, refer to the section entitled "SPRAYING INSTRUCTIONS - GROUND APPLICATION" which precedes this section.
2. Water Volume - Aerial application: Minimum of 30 litres per hectare when applied alone or when tank mixed with Buctril M.
3. When applying by aircraft, applicators are required to use spray nozzles or nozzle configurations and spraying pressures which will provide a COARSE droplet size distribution, with a volume medium diameter (VMD) greater than 350 microns.
4. Aerial drift is increased under certain meteorological conditions. Do not apply during periods of dead calm, when winds are gusty or when wind speed is greater than 15 km/hr.
5. For the protection of non-target habitats, overspray or drift to sensitive habitats must be avoided. A buffer zone of 76 metres is required between the downwind edge of the boom and sensitive terrestrial habitats including grasslands, forested areas, shelter belts, woodlots, hedgerows and shrublands. A buffer zone of 72 meters is required between the downwind edge of the boom and sensitive aquatic habitats such as sloughs, ponds, prairie potholes, lakes, rivers, streams and irrigation ditches, and the vegetation surrounding wetlands. Do not contaminate these habitats when cleaning and rinsing spray equipment or containers.
6. When a tank mixture is used, consult the label of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.
7. Ensure uniform application. To avoid uneven or overlapped application, use appropriate marking devices. Do not use human flaggers.

SPRAYER CLEAN-UP:

1. Thoroughly clean application equipment immediately after spraying. Ensure that all traces of the product are removed. The following recommendations are provided:
2. Drain and flush tank walls, boom and all hoses for ten minutes with clean water. Do not clean the sprayer near desirable vegetation, wells, or other water sources.
3. Remove the nozzles and screens and wash separately.
4. Dispose of all rinsings in accordance with provincial regulations.
5. If a broadleaf tank-mix partner is used, always check tank-mix partner label for any additional clean up procedures.

Resistance-Management Recommendations

For resistance management, Foothills A Herbicide is a Group 1 herbicide. Any weed population may contain or develop plants naturally resistant to Foothills A Herbicide and other Group 1 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field.

Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

Where possible, rotate the use of Foothills A Herbicide or other Group 1 herbicides with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group when such use is permitted.

Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.

Monitor treated weed populations for resistance development.

Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact company representatives at 1-877-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.

NOTE TO USER:

READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the use described below were developed by persons other than Syngenta Crop Protection Canada, Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Syngenta Crop Protection Canada, Inc. itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used on the crop listed below.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold Syngenta Crop Protection Canada, Inc. harmless, from any claims based on efficacy and/or phytotoxicity in connection with the use described below.

DIRECTIONS FOR USE IN THE OKANAGAN AND CRESTON FLATS REGION OF BC:

FOOTHILLS A Herbicide can be used for the control of wild oats, green foxtail and yellow foxtail in spring wheat and durum wheat in the Okanagan and Creston Flats Regions of British Columbia. For information on crop and weed stages, rates of application, mixing and spraying instructions and precautions see the appropriate sections elsewhere on this label.

Product names marked ® or ™ are registered trademarks of a Syngenta Group Company

† All other products listed are trademarks of their respective companies.

It must be noted FOOTHILLS™ A and FOOTHILLS™ B are the trademarks of Viterra. Viterra is a licensed user.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.