

PMRA Approved PH  
2009-4782  
22-JAN-2010

[LABEL]

GROUP	2	HERBICIDE
-------	---	-----------

**FirstStep™**

HERBICIDE

DRY FLOWABLE

For Sale for Use on Wheat (spring and durum), Barley and Summerfallow in the Prairie Provinces and the Peace River Region of British Columbia

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

REGISTRATION NO.: 29569 PEST CONTROL PRODUCTS ACT

GUARANTEE: Tribenuron-methyl 75%

WARNING – EYE IRRITANT  
POTENTIAL SKIN SENSITIZER  
Contains the allergens sulfites and milk

NET CONTENTS: 160 grams – 13.6 Kg

E. I. du Pont Canada Company  
Agricultural Products  
Box 2300 Streetsville  
Mississauga, Ontario  
L5M 2J4  
1-800-667-3925

Distributed by: Viterra Inc.  
2625 Victoria Avenue  
Regina, SK  
S4T 7T9

PRECAUTIONS:      WARNING - EYE IRRITANT.  
POTENTIAL SKIN SENSITIZER.  
DO NOT APPLY BY AIR.  
KEEP OUT OF REACH OF CHILDREN.  
MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN.  
AVOID BREATHING SPRAY MIST. USE WITH ADEQUATE VENTILATION.  
AVOID CONTACT WITH SKIN, EYES AND CLOTHING.  
DO NOT CONTAMINATE ANY BODY OF WATER.  
WEAR SUITABLE CHEMICAL RESISTANT GLOVES AND GOGGLES OR FACE SHIELD DURING MIXING AND LOADING.  
MAY IRRITATE SKIN AND EYES. WHEN USING, DO NOT EAT, DRINK OR SMOKE.  
WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING.  
REMOVE CONTAMINATED CLOTHING IMMEDIATELY AFTER USE.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: [www.croplife.ca](http://www.croplife.ca).

#### **FIRST AID**

In case of poisoning, contact a physician or poison control centre.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INGESTED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-441-3637 (24 hours).

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

#### **STORAGE AND DISPOSAL**

**STORAGE:** Store product in original container in a secure, dry area, away from fertilizer, seed, food or feed. Not for use or storage in or around the home. Keep container tightly closed.

**DISPOSAL:**

**FOR REFILLABLE CONTAINERS:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

#### **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the PEST CONTROL PRODUCTS ACT to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

®/™ Registered trademarks/trademarks of E. I. du Pont de Nemours and Company. E. I. du Pont Canada Company is a licensed user. Member of CropLife.

FIRSTSTEP™ is a trademark of Viterro Inc.

[DFU]

GROUP	2	HERBICIDE
-------	---	-----------

# FirstStep™

HERBICIDE

DRY FLOWABLE

For Sale for Use on Wheat (spring and durum), Barley and Summerfallow in the Prairie Provinces and the Peace River Region of British Columbia

COMMERCIAL

READ THE LABEL AND THIS BOOKLET BEFORE USING

REGISTRATION NO.: 29569 PEST CONTROL PRODUCTS ACT

GUARANTEE: Tribenuron-methyl 75%

WARNING – EYE IRRITANT  
POTENTIAL SKIN SENSITIZER  
Contains the allergen sulfites and milk

E. I. du Pont Canada Company  
Agricultural Products  
Box 2300 Streetsville  
Mississauga, Ontario  
L5M 2J4  
1-800-667-3925

Distributed by: Viterra Inc.  
2625 Victoria Avenue  
Regina, SK  
S4T 7T9

**PRECAUTIONS:**           WARNING - EYE IRRITANT.  
                                  POTENTIAL SKIN SENSITIZER.  
                                  DO NOT APPLY BY AIR.  
                                  KEEP OUT OF REACH OF CHILDREN.  
                                  MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH  
                                  SKIN. AVOID BREATHING SPRAY MIST. USE WITH ADEQUATE  
                                  VENTILATION.  
                                  AVOID CONTACT WITH SKIN, EYES AND CLOTHING.  
                                  DO NOT CONTAMINATE ANY BODY OF WATER.  
                                  WEAR SUITABLE CHEMICAL RESISTANT GLOVES AND GOGGLES OR  
                                  FACE SHIELD DURING MIXING AND LOADING.  
                                  MAY IRRITATE SKIN AND EYES. WHEN USING, DO NOT EAT, DRINK OR  
                                  SMOKE.  
                                  WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING.  
                                  REMOVE CONTAMINATED CLOTHING IMMEDIATELY AFTER USE.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: [www.croplife.ca](http://www.croplife.ca).

**FIRST AID:**

In case of poisoning, contact a physician or poison control centre IMMEDIATELY

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INGESTED:**            Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:**    Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-661-8323 (24 hours).

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

**IMPORTANT INFORMATION - READ BEFORE USING**

Injury to or loss of desirable trees, shrubs, flowers, vegetables, fruits or other vegetation may result from failure to observe the following: Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on crops.

**DO NOT APPLY BY AIR.**

Do not apply in high winds or on hot days (over 27 degrees C).

Carefully observe sprayer clean-up instructions, as spray tank residue may damage crops other than spring wheat, durum wheat, barley or summerfallow. Thoroughly clean all traces of tank mix from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately). Wash away any spray mixture from the outside of spray tank, nozzle or spray rig.

A MINIMUM RECROPPING INTERVAL OF 2 MONTHS SHOULD BE LEFT BETWEEN THE APPLICATION OF THIS PRODUCT AND SEEDING OF THE NEXT CROP. After 2 months, the following crops can be seeded: canola, flax, lentils, and alfalfa.

DO NOT APPLY FIRSTSTEP WITH A CARRIER OTHER THAN WATER OR ANY ADDITIVE OTHER THAN THOSE INDICATED ON THIS LABEL. Other carriers may accelerate breakdown of this product and reduce its effectiveness.

### **GENERAL INFORMATION**

FirstStep provides control of certain broad-leaved weeds in spring planted wheat (including durum), barley and summerfallow.

Warm, moist growing conditions promote active weed growth and enhance the activity of FirstStep by allowing maximum foliar uptake and contact activity. Weeds hardened off by environmental stress such as cold weather, drought stress or excessive heat may not be adequately controlled or suppressed, and regrowth may occur. FirstStep may only be applied using ground spray equipment. For best results, ensure thorough spray coverage of target weeds.

FirstStep rapidly stops growth of susceptible weeds. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending on growing conditions and weed susceptibility. Degree of control and duration of effect depend on weed sensitivity, weed size, spray coverage and growing conditions. Activity of the herbicide mixture may be delayed by cold, dry conditions after application.

### **DIRECTIONS FOR USE:**

#### **FOR CONTROL OF FLIXWEED AND STINKWEED (FALL ROSETTES AND SPRING SEEDLINGS) IN SUMMERFALLOW**

FirstStep + 2,4-D ESTER TANK MIX:

Apply FirstStep at 10 g/ha with 2,4-D Ester at 420 g/ha (0.6 L 2,4-D Ester LV-700). AGRAL 90 is not required in this tank mix.

Apply the tank mix in the spring after emergence, up to the early flowering stage of the target weeds. Only weeds that have emerged at the time of application will be controlled.

#### **CONTROL OF GRASS AND BROADLEAF WEEDS IN SUMMERFALLOW**

FirstStep + ROUNDUP\* TRANSORB, ROUNDUP\* DRY or TOUCHDOWN\* 600

Apply FirstStep at 10 g/ha with ROUNDUP\* TRANSORB at 1.25 L/ha, ROUNDUP\* DRY at 0.70 kg/ha or TOUCHDOWN\* 600 at 1.0 L/ha in a total spray volume of 55 - 110 L/ha. Refer to the following table for the weeds controlled and application timing.

<b>Weeds Controlled</b>	<b>Growth Stage</b>
Canada fleabane	Up to 8 cm
Canada thistle (suppression)	rosette
common ragweed	Up to 8 cm
cow cockle	Up to 3 leaf
dandelion	Up to 15 cm
downy brome	Up to 15 cm
flixweed	Up to 15 cm
giant foxtail	Up to 15 cm
green foxtail	Up to 15 cm
hempnettle	Up to 15 cm
kochia	Up to 15 cm
lady's thumb	Up to 15 cm
lamb's quarters	Up to 15 cm
narrow-leaved hawk's beard	Up to 8 cm
Persian darnel	Up to 15 cm
redroot pigweed	Up to 15 cm
Russian thistle	Up to 15 cm
stinkweed	Up to 15 cm
volunteer barley	Up to 15 cm
volunteer canola	Up to 15 cm
volunteer flax	Up to 15 cm
volunteer wheat	Up to 15 cm
wild buckwheat	Up to 3 leaf
wild mustard	Up to 15 cm
wild oats	Up to 15 cm

Allow at least 10 days to elapse before tillage.

**FOR PRE-SEED WEED CONTROL IN SPRING WHEAT (INCLUDING DURUM) AND BARLEY AND WEED CONTROL IN SUMMERFALLOW**

FirstStep + ROUNDUP\* ORIGINAL OR ROUNDUP\* TRANSORB

Apply 10 grams of FirstStep per hectare, tank mixed with Round-Up\* Original or Roundup\* Transorb at 0.75 – 1.0L/Ha, with Agral\* 90 at 0.35% v/v (3.5 L/1000 L) in a total spray volume of 55-110 L/ha.

In summerfallow, allow at least 10 days to elapse before tillage.

#### WEEDS CONTROLLED

Dandelion – top growth control  
Canola, including glyphosate tolerant canola (e.g. Roundup Ready\* canola)  
Green foxtail  
Kochia  
Lady's-thumb  
Lamb's-quarters  
Narrow-leaved hawk's beard  
Stinkweed  
Volunteer barley  
Volunteer wheat  
Wild buckwheat  
Wild mustard  
Wild oats

#### WEEDS SUPRESSED

Canada thistle\*  
Redroot pigweed\*  
Russian thistle\*

\*Weed suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

## APPLICATION RATES AND TIMING

Tank-mix with FirstStep at 10 g/Ha (7.5 g ai/ha)		
Roundup* Application Rate	Weed	Application Stage
Roundup Original* or Roundup Transorb* at 0.75 L/Ha	Green Foxtail	Up to 8 cm
	Kochia	Up to 10 cm
	Lady's-thumb	Up to 8 cm
	Lamb's-quarters	Up to 10 cm
	Narrow-leaved hawk's beard (suppression)	Rosettes and seedlings
	Redroot Pigweed (suppression)	Up to 10 cm
	Russian Thistle (suppression)	Up to 10 cm
	Stinkweed	Up to 8 cm
	Volunteer Barley	Up to 8 cm
	Volunteer Canola (including glyphosate tolerant)	Any stage
	Volunteer Wheat	Up to 8 cm
	Wild Buckwheat	1-3 leaf
	Wild mustard	Up to 8 cm
	Wild oats	Up to 8 cm (1-3 leaf)
Roundup Original* or Roundup Transorb* at 1.0 L/Ha	Canada Thistle (suppression of top growth)	Rosettes
	Dandelion (top growth control)	Seedlings, non-flowering rosettes

## SEEDING OF CROP AFTER APPLICATION

Fields treated with FirstStep + ROUNDUP\* ORIGINAL or ROUNDUP TRANSORB can be seeded with spring wheat (including durum) or barley a minimum of 24 hours after application.

## MIXING INSTRUCTIONS

1. Add one-third of the necessary volume of water to the spray tank.
2. With the agitator running, add the required amount of FirstStep.
3. When FirstStep is in suspension, add the required amount of the appropriate tank mix partners.

FirstStep + 2,4-D Ester: once FirstStep is completely in suspension, add the required amount of 2,4-D Ester while maintaining agitation. Do not add a surfactant.

FirstStep + ROUNDUP TRANSORB\* or ROUNDUP\* DRY or TOUCHDOWN\* 600: once FirstStep is completely in suspension, add the required amount of Roundup Transorb\*, Roundup\* Dry or Touchdown\* 600 while maintaining agitation.

4. Add the rest of the water. If an antifoam agent is required, add this last.

5. Use a minimum spray volume of 55 L/ha on spring wheat (including durum), barley and summerfallow.
6. For repeat tankloads, reduce the material remaining in the tank to 10% of the original volume or less before proceeding with step 1. If this is not possible, pre-slurry FirstStep in a small amount (5-10 L) of water before adding to the tank.

Continuous agitation is required to keep FirstStep in suspension. If the spray preparation is left standing without agitation, thoroughly agitate to resuspend the mixture before spraying. Use spray preparation within 24 hours or product degradation may occur.

### **MINIMUM RECROPPING INTERVAL AND CROP ROTATION GUIDELINES**

A minimum recrop interval of 2 months should be left between the application of this product and seeding of the next crop. The following crops can be seeded two months after application of FirstStep + 2,4-D Ester: canola, flax, lentils, and alfalfa.

### **EQUIPMENT-SPRAY VOLUMES**

Apply the spray mixture uniformly with properly calibrated ground equipment only. Ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended (minimum 55 litres spray volume per hectare). Use 50 mesh filter screens or larger (metal or nylon).

Continuous agitation is required to keep the spray materials in suspension. Apply uniformly and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

Reduce drift with high water volume, low pressure, coarse sprays and drop nozzles. Do not apply with more than 275 kPa spray pressure.

DO NOT APPLY BY AIR.

### **SPRAYER CLEANUP**

It is difficult to remove all traces of 2,4-D Ester and MCPA Amine from a sprayer. A sprayer used to apply these herbicides should not be used to spray crops sensitive to these herbicides. The following clean-out procedure will remove traces of FirstStep from the sprayer.

1. Drain tank; then flush tank, boom and hoses with clean water for a minimum of ten minutes to remove all visible residues.
2. Fill the tank with clean water then add 1 litre household AMMONIA (containing a minimum of 3% ammonia) per 100 litres of water. Fill boom and hoses with solution and allow sprayer to sit for 15 minutes. Drain.
3. Repeat step 2.
4. Nozzles and screens should be removed and cleaned separately. To remove traces of ammonia, rinse the tank, hoses and booms thoroughly with clean water.
5. Dispose of tank rinseate according to Provincial directions.

CAUTION: Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour that can cause eye, nose, throat, and lung irritation. Do not clean equipment in an enclosed area.

### **RESISTANCE-MANAGEMENT RECOMMENDATIONS**

For resistance management, FirstStep is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to FirstStep and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the used of FirstStep or other Group 2 herbicides with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted
- ◆ Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- ◆ Monitor treated weed populations for resistance development.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local DuPont representative or the DuPont hotline at 1-800-667-3925 for further information.

#### **USE PRECAUTIONS:**

Licensed wheat and barley varieties have shown acceptable tolerance to FirstStep. Consult DuPont prior to use on unlicensed varieties for crop tolerance information.

Do not apply to irrigated land where tail water will be used to irrigate other cropland.

Do not contaminate irrigation water.

Do not use on highly variable soils that have large gravelly or sandy areas, eroded knolls or calcium deposits.

If rain occurs soon after application, control may be reduced. At least 4-6 hours of dry weather is needed to allow the tank mixture to be absorbed by weed foliage. Environmental conditions that slow the drying of the spray mixture on the foliage such as high relative humidity, cool air temperatures or cloud cover, may increase the time required.

Use a maximum of 10 grams of FirstStep per hectare per year on spring planted wheat, barley, and summerfallow.

FirstStep applied to summerfallow, spring wheat, durum wheat or barley that is stressed by severe conditions such as drought, low fertility, saline soils, waterlogged soils (soils at or near field capacity), disease or insect damage may result in crop injury. Drought, disease or insect damage following application may also result in crop injury, grade or yield loss.

Do not apply to spring wheat, durum wheat or barley undersown with legumes or grasses as injury to the forage may result.

Do not remove soil from treated fields for use on lawns, in gardens or for backfill.

Under certain conditions such as prolonged cool weather temperature or wide fluctuations in day/night temperatures just prior to or soon after treatment, temporary yellowing and/or crop stunting may occur.

OVERSPRAY OR DRIFT TO IMPORTANT WILDLIFE HABITATS SUCH AS SHELTERBELTS, WETLANDS, SLOUGHS OR DRY SLOUGH BORDERS, OR WOODLOTS, SHOULD BE AVOIDED. LEAVE A 15-METER BUFFER ZONE BETWEEN THE LAST SPRAY SWATH AND THE EDGE OF ANY OF THESE HABITATS.

Control of weeds growing in wheel tracks may be reduced if FirstStep is applied under dry, dusty conditions.

## **STORAGE AND DISPOSAL**

### **STORAGE:**

Store products in original containers only, away from fertilizer, seeds, food or feed.

Not for use or storage in or around the home. Keep container closed.

### **DISPOSAL:**

#### **FOR REFILLABLE CONTAINERS:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

®/™ Registered trademarks/trademarks of E. I. du Pont de Nemours and Company. E. I. du Pont Canada Company is a licensed user. Member of CropLife.

FIRSTSTEP™ is a trademark of Viterro Inc.

\* All other products mentioned are trademarks of their respective companies.