

2008-5735
06-FEB-2009

(Front Panel)

GROUP HERBICIDE

VITERRA

2,4-D Amine 600

LIQUID HERBICIDE

AGRICULTURAL

READ THE LABEL AND THE BOOKLET BEFORE USING.

GUARANTEE: 2,4-D (Present as Dimethylamine salt): 560 g a.e./L
Warning, contains the allergen soy.

REGISTRATION NO: 29248
PEST CONTROL PRODUCTS ACT

Net Contents: **10 litres**

WARNING



POISON

Manufactured for:

Viterra Inc.
2625 Victoria Avenue
Regina, SK S4T 7T9

For Emergencies Call: 1-877-943-2876

By:

Interprovincial Cooperative Limited,
P.O. Box 1050, Saskatoon, Saskatchewan S7K 3M9
For Technical Support Call: 1-204-233-3461

VITERRA is a trademark of Viterra Inc.

RC 000-1208

DETACH LABEL AT PERFORATION. THIS PORTION MUST REMAIN ATTACHED TO THE CONTAINER.

PRECAUTIONS: Handling precautions: KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes, skin and clothing. May cause skin irritation. Avoid inhaling vapours or mist. **Protective Clothing and Equipment: Handling the concentrate (mixing and loading):** wear a long-sleeved shirt, long pants, coveralls, chemical-resistant gloves, socks and shoes. Rinse gloves before removal. Coveralls or a chemical resistant apron should be worn when pouring from containers greater than 5 Litres.

Handling the dilute spray solution (during application or repairing or cleaning equipment): Wear a long-sleeved shirt, long pants, coveralls, chemical-resistant gloves, socks and shoes. Rinse gloves before removal. Gloves are not required during application when applicator is in an enclosed tractor or an enclosed airplane cockpit. **Operator use precautions:** Wear freshly laundered clothing and clean protective equipment daily. Rinse gloves before removal. Wash hands before eating, drinking, using tobacco or using the toilet. If herbicide penetrates clothing remove immediately, then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. After using this product, remove clothing and launder separately and promptly, and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry. After work, remove all clothing and shower using soap and water. **GENERAL PRECAUTIONS:** Handle with care and mix only in a closed container. Use only in a well ventilated area. May cause damage to susceptible crops, ornamentals and other plants, in minute quantities. Avoid drift to desirable vegetation. Do not spray during periods of high wind. Coarse sprays are less likely to drift. Do not contaminate any body of water. Avoid contamination of foods. Keep in original container during storage. Do not re-use container unless otherwise specified. Store the container tightly closed away from seeds, fertilizers, plants and foodstuffs. Store above 0 degrees C. **SHAKE WELL BEFORE USING.** Clean spray equipment thoroughly after use. If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at: www.croplife.ca.

Environmental precautions: DO NOT apply this product directly to freshwater habitats such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands, estuaries or marine habitats. DO NOT contaminate irrigation/drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes. **TOXIC** to broadleaf terrestrial plants. This product may harm other broadleaf plants in the vicinity of the treatment area. Observe buffer zones specified under DIRECTIONS FOR USE. **LEACHING:** To minimize possible contamination of groundwater, the use of spot treatment application is recommended in areas where soils are permeable (e.g., sandy soil) or the water table is shallow. **RUN-OFF:** To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, bare soil, compacted soil, or clay. Avoid application of this product when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a strip of untreated vegetation between the treated area and the edge of the water body.

FIRST AID: in case of poisoning, call a physician or poison control centre IMMEDIATELY. **If on skin or clothing,** take off contaminated clothing. Rinse skin IMMEDIATELY with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. **If in eyes,** hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If swallowed,** call a poison control centre or doctor IMMEDIATELY for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **If inhaled,** move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. Take the container, label, or product name and Pest Control Product registration number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treatment of any systemic intoxication should be primarily symptomatic and supportive. This product may cause severe irritation to the eyes. Overexposure to 2,4-D may cause coughing, burning, dizziness, or temporary loss of muscle coordination. Other possible effects of overexposure include fatigue, muscle weakness and nausea.

RINSING AND DISPOSAL INFORMATION: For Recyclable Containers: Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site: 1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. 2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. **For Returnable Containers:** Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer). **For Returnable/Refillable Containers:** For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

(Detachable booklet)

DETACH LABEL INSTRUCTIONS AT PERFORATION

GROUP 4 HERBICIDE

VITERRA

2,4-D Amine 600

LIQUID HERBICIDE

AGRICULTURAL

READ THE LABEL AND THE BOOKLET BEFORE USING.

GUARANTEE: 2,4-D (Present as Dimethylamine salt): 560 g a.e./L
Warning, contains the allergen soy.

REGISTRATION NO: 29248
PEST CONTROL PRODUCTS ACT

Net Contents: **10 litres**



WARNING

POISON

Manufactured for:

Viterra Inc.
2625 Victoria Avenue
Regina, SK S4T 7T9

For Emergencies Call: 1-877-943-2876

By:

Interprovincial Cooperative Limited,
P.O. Box 1050, Saskatoon, Saskatchewan S7K 3M9
For Technical Support Call: 1-204-233-3461

VITERRA is a trademark of Viterra Inc.

RC 000-1208

LABEL INDEX – VITERRA 2,4-D Amine 600 Liquid Herbicide

Spring Wheat, Barley, Rye	GO TO SECTION 1	Strawberries (Eastern Canada Only)	GO TO SECTION 12
Winter Wheat, Fall Rye	GO TO SECTION 2	Raspberries (Eastern Canada Only)	GO TO SECTION 13
Field Corn	GO TO SECTION 3	Cranberries	GO TO SECTION 14
Weeds Controlled/Rates of Application	GO TO SECTION 4	Bearing Fruit Trees: Apple, Peach, Pear, Plum, Apricot, Sweet and Sour Cherry	GO TO SECTION 15
Forage Sorghum, Forage Millet	GO TO SECTION 5	Weeds and Brush On Non-Crop / Non-Grazed Land	GO TO SECTION 16
Established Pastures, Rangeland (Without Legumes)	GO TO SECTION 6	Tank Mixes	GO TO SECTION 17
Grass Crops for Forage and Seed Production	GO TO SECTION 7	Mixing and Application	GO TO SECTION 18
Stubble Land, Roadsides, and Uncropped Land	GO TO SECTION 8	Precautions	GO TO SECTION 19
Pre-Harvest / Grazing Intervals	GO TO SECTION 9	First Aid	GO TO SECTION 20
Turf, Established Fairways, Golf Courses, Lawns, Parks	GO TO SECTION 10	Toxicological Information	GO TO SECTION 21
Asparagus (Eastern Canada Only)	GO TO SECTION 11	Rinsing and Disposal Information	GO TO SECTION 22

DIRECTIONS FOR USE:

SECTION 1 - SPRING WHEAT, BARLEY, RYE

Treatment Stage: Ground or aerial application. Treat from the 4th leaf expanded stage to the early flag leaf (shot blade) stage. (Do not treat crop underseeded to legumes.)

GO TO SECTION 4

SECTION 2 - WINTER WHEAT, FALL RYE

Treatment Stage: Ground or aerial application. Treat in the spring from full tillering until just before the flag leaf (shot blade) stage. Do not apply to seedling winter cereals in the fall.

GO TO SECTION 4

SECTION 3 - FIELD CORN

Treatment Stage: Ground application only. As an overall spray, before the corn is 15 cm tall (leaf extended) and/or before the 6-leaf stage. Application as an overall spray at later stages will damage the corn. For later applications (corn 15-75 cm, leaf extended), use a shielded spray or a directed spray with drop pipes. Keep spray off corn foliage. Do not apply within 2 weeks of silking and tasselling. Avoid cultivation for 2 weeks following spraying. Use at least 100L/ha water. Do not add oil or surfactant.

GO TO SECTION 4

SECTION 4 - WEEDS CONTROLLED AND RATES OF APPLICATION

Susceptible Weeds: Annual sow-thistle, bluebur (up to 4 leaf stage), burdock (before the 4-leaf stage), cocklebur, common ragweed, daisy fleabane, false flax, false ragweed, flixweed, goat's-beard, giant ragweed, kochia, lamb's-quarters, mustards (except dog and tansy), narrow-leaved hawk's-beard (susceptible in the fall and at 1-2 leaf stage in the spring), common plantain, prickly lettuce, redroot pigweed, russian pigweed, russian thistle, shepherd's-purse, stinging nettle, stinkweed, sweet clover (seedling), thyme-leaved spurge, volunteer canola, wild radish, wild (prairie) sunflower.

Optimum Stage of Weed Growth: 2 to 4 leaf, growing actively.*

Rates to Use on Susceptible Weeds:

Small seedlings, good growing conditions:	600 mL/ha
Weeds in bud, dry/cold conditions, or heavy infestation:	950 mL/ha

* Weed control may be reduced when applied beyond optimum stage of weed growth.

Harder to Control Weeds: Blue lettuce, biennial wormwood, common purslane, curled dock (up to 4 leaf), dandelion in lawn, dog and tansy mustard, flixweed(spring prior to bolting), groundsel, gumweed, hairy galinsoga, hawkweed, heal-all, hedge bindweed, knotweed(up to 4 leaf), narrow-leaved hawk's-beard (spring prior to bolting), oak-leaved goosefoot, peppergrass, pineappleweed, prostrate pigweed, Russian knapweed, sheep sorrel, tumble pigweed, velvet leaf, yellow rocket (before the 4 leaf stage).

Optimum Stage of Weed Growth: 2-4 leaf , growing actively.*

Rates to Use on Harder to Control Weeds:

Small seedlings, good growing conditions:	1.0 L/ha
Weeds in bud, dry/ cold conditions, or heavy infestations:	1.5 L/ha

NOTE: These high rates may result in deformities in the crop, but improved weed control should result in higher yields.

* Weed control may be reduced when applied beyond optimum stage of weed growth.

TOP GROWTH CONTROL ONLY:

Biennial wormwood, blue lettuce, bull thistle, burdock, buttercup, Canada thistle, field dandelion, field bindweed, gumweed, hedge bindweed, hoary cress, leafy spurge, mouse-eared chickweed*, perennial sow-thistle, Russian knapweed, tartary buckwheat, teasel, volunteer sunflower, yellow rocket (controlled with applications before the 4-leaf stage).

- Use highest rate listed below for suppression only.

Rates of Application Top Growth Control Only:

Small seedlings, good growing conditions:	1.0 L/ha
Large weeds, dry/cold conditions, or heavy infestations:	1.5 L/ha

NOTE: For use in-crop. These rates may result in significant injury on some crops and should only be used where some crop injury can be tolerated.

Maximum Safe Rates of Application For Crops:

Spring wheat, barley, rye:	1.0 L/ha
Winter wheat, fall rye:	900 mL/ha
Field corn:	1.0 L/ha

Do not exceed these maximum rates in the weed control tables in any crop.

GO TO SECTION 9

SECTION 5 - FORAGE SORGHUM, FORAGE MILLET

Treatment Stage: Ground application only. Apply post emergent to the weeds and crop when the crop is at the 4-6 leaf stage, before closure of the canopy.

Susceptible Weeds: As listed in Section 4.

Rates:	Small seedlings, good growing condition:	500 mL/ha
	Weeds in bud, dry conditions, very heavy infestations:	1.0 L/ha

Harder to Kill Weeds: As listed in Section 4.

Rate:	Small seedlings, good growing conditions:	1.0 L/ha
--------------	---	----------

GO TO SECTION 9

SECTION 6 - ESTABLISHED PASTURES, RANGELAND (WITHOUT LEGUMES)

Treatment Stage: Ground or aerial application. Apply at a time of rapid growth, usually May, June and/or September. Apply after pasture has been grazed or cut and weed growth has resumed. A second treatment may be required. Do not apply in freshly seeded grasses and pastures until well established. Under conditions of good growth and adequate soil moisture, sprays can be applied up to 2 weeks before normal frost time. Plants poisonous to livestock (eg. tansy ragwort, water hemlock, poison hemlock, death camas) may appear more palatable to grazing livestock and could be preferentially grazed by livestock after application of VITERRA 2,4-D Amine 600 Liquid Herbicide. Livestock should not be allowed to graze treated areas of such plant growth until poisonous plants have been removed, are completely dead, or have resumed normal growth habit.

Weeds Controlled: As listed in Section 4

Rates to Use: Up to 1.5 to 4.0 L/ha in 100-300 litres of water per hectare.
Use the higher rate for perennial weeds.

GO TO SECTION 9

SECTION 7 - GRASS CROPS FOR FORAGE AND SEED PRODUCTION

Treatment Stage: Ground application only.

Seedling Grasses-Stand Establishment – For seedling grasses apply from the 3 leaf to just before the shot blade stage of development.

Established Grasses: Apply in the spring up to the shot-blade stage of the grasses or in the fall after harvest to control annual and biennial weed species.

Weeds Controlled: As listed in Section 4

Rates to Use: **Seedling Grasses-Stand Establishment:**
0.49 to 0.95 L/ha of product in 100 to 200 L/ha of water

Established grasses:

0.6–1.9 L/ha in 100-300 L/ha of water.

On established grass seed crops, do not exceed 0.95 L/ha. Do not apply during flower development or pollination as seed yields may be reduced.

Late germinating weeds such as red bartsias may be controlled at 0.95 L/ha of product after the first hay crop is removed.

GO TO SECTION 9

SECTION 8 - STUBBLE LAND, ROADSIDES, AND UNCROPPED LAND.

Treatment Stage: Ground or aerial application for stubble land and uncropped land. **DO NOT APPLY BY AIR TO ROADSIDES.**

Susceptible Weeds: As listed in Section 4.

Rates of Use on Susceptible Weeds:

Seedling Weeds:	1.50 L/ha
Weeds in bud, dry/cool conditions, heavy infestations:	2.75 L/ha

Harder to Control Weeds: As listed in Section 4.

Rate to Use on Harder to Control Weeds:

Seedling Weeds (up to 4 leaf):	2.75 L/ha
Weeds in bud, dry/cool conditions, heavy infestations, perennials and small brush:	4 L/ha

NOTES:

Spray thoroughly using 100 litres or more of water per hectare by ground. Use the higher rate for perennial weeds and susceptible woody growth. A second treatment may be required. Cultivating grain stubble and spraying after regrowth may aid in control. Do not apply to freshly seeded grasses until well established. Under conditions of good growth and adequate soil moisture, sprays can be applied up to 2 weeks before normal frost time. Plants poisonous to livestock (eg. tansy ragwort, water hemlock, poison hemlock, death camas) may appear more palatable to grazing livestock and could be preferentially grazed by livestock after application of VITERRA 2,4-D Amine 600 Liquid Herbicide. Livestock should not be allowed to graze treated areas of such plant growth until poisonous plants have been removed, are completely dead, or have resumed normal growth habit.

GO TO SECTION 9

SECTION 9 - PRE-HARVEST/GRAZING INTERVALS (cereal crops, field corn, forage sorghum, forage millet, established pastures and rangeland, grass crops for forage).

1. Do not permit lactating dairy animals to graze fields within 7 days after application.
2. Do not harvest forage or cut hay within 30 days after application.
3. Withdraw meat animals from treated fields at least 3 days before slaughter.

GO TO SECTION 18

SECTION 10 - TURF, ESTABLISHED FAIRWAYS, GOLF COURSES, LAWNS, PARKS.

Treatment Stage: DO NOT apply by air. Treat in Spring (May & early June) and early September before grasses are in the flag leaf (shot blade) stage for the best results.

Weeds Controlled: Plantain, Dandelion, Curled Dock, Mustard, Pigweed, Ragweed, Shepherd's Purse and other weeds susceptible to 2,4-D (see section 4 for a complete listing of weeds controlled)

Rate to Use: 1.9 L/ha in 100 to 200 L/ha of spray solution.

Maximum Application Frequency to Turf:

DO NOT apply more than two broadcast applications per season. This does not include spot treatments.

Re-entry Interval for Turf:

Do not allow people (other than applicators) or pets on treatment area during application. Do not enter treated areas until spray has thoroughly dried.

Application notes: Do not apply to bentgrass or newly seeded lawns/turf as injury may occur. Avoid application if rain is expected within 24 hours or if weather is hot (above 27 deg. C) and/or dry. Avoid mowing within 4 days before and after application. If weed populations do not warrant a broadcast application (e.g. entire lawn), consider spot treatments that target only weedy areas. This product is only effective when applied to the leaves of actively growing weeds. This product will not prevent new weeds – apply only when weeds are present. DO NOT apply to the exposed roots of trees and ornamentals.

Field Sprayer Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification.

Buffer Zones

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats such as grasslands, forested areas, shelter belts, woodlots, hedgerows, pastures, rangelands and shrublands. The buffer zones in the table below are reduced by 70% with the use of shrouds and 30% with the use of cones on field sprayers. A further substantial reduction can be achieved by using ASAE coarse droplet size instead of medium droplet size.

When a tank mixture is used, consult the labels of the tank-mix partners and use the largest (most restrictive) buffer zone recommended for any of the products.

	Buffer Zones(meters)		
	Using Tractor Pulled Boom Sprayer		
	Unshielded Boom (Buffer Zone)	Shrouded Boom (Shrouds)	Coned Boom (Cones)
Medium Droplet Size 250-350 um-vmd	20	6	14
Coarse Droplet Size 350-450 um-vmd	4	1	3

vmd= volume mean diameter.

GO TO SECTION 18

SECTION 11 – ASPARAGUS (EASTERN CANADA ONLY)

Treatment stage: Do not apply by air. Apply early in the season immediately after a thorough cutting or disking. Application can also be made 3 weeks after a post harvest disking using a directed spray to avoid the ferns.

Susceptible weeds: See Section 4 for weeds controlled.

Rates of application: 1.7 to 2.3 litres per hectare. Use the lower rate for weeds identified as susceptible weeds and the higher rate for weeds identified as harder to control or where top growth control is acceptable as under section 4. Apply in 50 to 200 litres of spray solution per hectare.

Notes: If emerged spears are contacted directly by spray or drift, they may become distorted or twisted. Spears emerging after application will be normal.

GO TO SECTION 18

SECTION 12 – STRAWBERRIES (EASTERN CANADA ONLY)

Treatment methods: Do not apply by air.

After Planting: Apply 2 to 4 weeks AFTER PLANTING but before the runners root in June. A low pressure nozzle and coarse spray is recommended.

Stand Renovation: Apply after harvest but before mowing. Delay mowing a few days after application. DO NOT apply between mid-August and fall dormancy due to damage to flower bud formation. Veestar is more sensitive than Redcoat. This treatment may cause distortion to the leaves.

Rates of application: After planting: 0.95 L/ha. **Stand renovation:** 1.7 L/ha
Apply in 100 to 200 L/ha of spray solution.

GO TO SECTION 18

SECTION 13 – RASPBERRIES – (Eastern Canada ONLY)

Treatment method: Do not apply by air.

Apply any time EXCEPT DURING BLOOM. Avoid spraying new shoots.

Rate of application: 0.95 L/ha. Apply in 100 to 200 L/ha of spray solution. Higher rates may be used for spot treatments.

GO TO SECTION 18

SECTION 14 – CRANBERRIES

Treatment method: Do not apply by air.

Apply as an over the top, wipe-on application to actively growing weeds (June and July). Injury will result if crop is contacted. Apply only one application per year and observe a pre-harvest interval of 50 days.

Rate of application: Apply at a dilution rate of 1 part of VITERRA 2,4-D Amine 600 Liquid Herbicide to 2.6 parts of water (27.5% solution).

Special weeds controlled: In addition to those noted in Section 4 of susceptible weeds, Hardhack, St. Johnswort, Alder and Purple Aster.

SECTION 15 -BEARING FRUIT TREES: Apple, Peach, Pear, Plum, Apricot, Sweet and Sour Cherry

Treatment Stage: Apply in the early spring after weed emergence, but in compliance with pre-harvest interval of 80 days or post-harvest to actively growing broadleaf weeds. DO NOT APPLY BY AIR.

Weeds Controlled: Plantains, Dandelion, Curled Dock, Mustards, Redroot Pigweed, Prostrate Pigweed, Russian Pigweed, Common Ragweed, Shepherd's Purse, Lamb's-Quarters, Stinkweed, Cocklebur, Prickly Lettuce, Narrow-leaved Hawk's Beard (fall rosette or spring at 1-2 leaf stage), Russian Thistle, seedling Sow Thistle and seedling Canada Thistle.

Rate of Application: 1.7 L/ha

Pre-Harvest Interval: Do not apply to weeds within 80 days of harvest of fruit crop.

Precaution: Do not apply when useable fruit is on the orchard floor.

Re-Entry Interval: Do not allow people (other than applicators) or pets into treatment area during application. Do not enter treated area until spray has thoroughly dried.

GO TO SECTION 18

SECTION 16 - WEEDS & BRUSH ON NON-CROP / NON-GRAZED LAND

Susceptible Species: Consult provincial spray calendars for a complete list of susceptible species. Ground or aerial application.

Rates to Use: 2.75 – 7.90 L/ha
Use the lower rate for species that are more susceptible such as chokecherry, western snowberry, willow and poplars.

Rate of Carrier: Use sufficient water to wet foliage thoroughly. 1000 to 3000 L/ha of spray solution may be required if applying by ground. Where only weeds occur, use 100 to 200 L/ha of spray solution. For aerial applications use 55 to 165 litres of spray solution per hectare.

Notes: Spray brush species up to 3 metres tall after foliage is well developed. Applications soon after leaves are fully open usually give the best results. Apply sufficient spray volume to thoroughly wet all plant parts to the ground line. Brush or trees above 3 m tall should be cut close to the ground and sprayed when the re-growth is over 0.6 m tall. With good growing conditions and adequate soil moisture, sprays may be applied up to 2 to 3 weeks before normal frost time.

GO TO SECTION 18

SECTION 17 – TANK MIXES

Use tank mixes only when they are recommended for use. Read and follow the directions, aerial application directions, warnings, precautions, and grazing limitations on both products used in tank-mix. When tank-mixing, observe the most stringent precautionary measures for handling and mixing stated on both product labels. See appropriate tank-mix product labels for full details of use patterns, rates and timings of applications and limits on application.

GO TO SECTION 18

SECTION 18 - MIXING AND APPLICATION

1. Fill tank 1/2 full with water. Start agitation.
2. Add VITERRA 2,4-D Amine 600 Liquid Herbicide.
3. When tank-mixes are used, consult the tank-mix product label for full mixing instructions.
4. Complete filling of tank.
5. Use 50-200 litres of water per hectare for ground application unless otherwise specified or on tank-mix product labels.
6. Do not exceed 275 kPa pressure for ground application

Use Precautions for Aerial Application (see specific sections for crops): Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment. Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions: Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application, as outlined in the *National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides*. Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed. Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelterbelt) or aquatic habitat.

Operator Precautions: Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of pre-mixed chemicals with a closed system is permitted. It is desirable that the pilot has communication capabilities at each treatment site at the time of application. The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls, and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

GO TO SECTION 19

SECTION 19 - PRECAUTIONS

Handling precautions: KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes, skin and clothing. May cause skin irritation. Avoid inhaling vapours or mist. **Protective Clothing and Equipment: Handling the concentrate (mixing and loading):** wear a long-sleeved shirt, long pants, coveralls, chemical-resistant gloves, socks and shoes. Rinse gloves before removal. Coveralls or a chemical resistant apron should be worn when pouring from containers greater than 5 Litres.

Handling the dilute spray solution (during application or repairing or cleaning equipment): wear a long-sleeved shirt, long pants, coveralls, chemical-resistant gloves, socks and shoes. Rinse gloves before removal. Gloves are not required during application when applicator is in an enclosed tractor or an enclosed airplane cockpit. **Operator use precautions:** Wear freshly laundered clothing and clean protective equipment daily. Rinse gloves before removal. Wash hands before eating, drinking, using tobacco or using the toilet. If herbicide penetrates clothing remove immediately, then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. After using this product, remove clothing and launder separately and promptly, and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry. After work, remove all clothing and shower using soap and water.

GENERAL PRECAUTIONS: Handle with care and mix only in a closed container. Use only in a well ventilated area. Avoid contamination of foods. Keep in original container during storage. Do not re-use container unless otherwise specified. Store the container tightly closed away from seeds, fertilizers, plants and foodstuffs. Store above 0 Deg. C. SHAKE WELL BEFORE USING. Clean spray equipment thoroughly after use. **Transfer System:** For containers larger than 20 L, use a transfer system that avoids open pouring when transferring the liquid concentrate from such containers into the spray tank. If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at: www.croplife.ca.

ENVIRONMENTAL PRECAUTIONS

DO NOT apply this product directly to freshwater habitats such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands, estuaries or marine habitats. DO NOT contaminate irrigation/drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Toxic to broadleaf terrestrial plants. This product may harm other broadleaf plants in the vicinity of the treatment area. Observe buffer zones specified under DIRECTIONS FOR USE ON TURF

Leaching

To minimize possible contamination of groundwater during application to turf areas, the use of spot treatment application is recommended in areas where soil is permeable (e.g, sandy soil) or the water table is shallow.

Run-off

To reduce runoff from treated turf areas into aquatic habitats, avoid application to areas with a moderate to steep slope, bare soil, compacted soil or clay.

Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a strip of untreated vegetation between the treated area and the edge of the water body.

RESISTANCE MANAGEMENT RECOMMENDATIONS: For resistance management, VITERRA 2,4-D Amine 600 Liquid Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to VITERRA 2,4-D Amine 600 Liquid Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance: Where possible, rotate the use of VITERRA 2,4-D Amine 600 Liquid Herbicide or other Group 4 herbicides with different herbicide groups that control the same weeds in a field. Use tank mixtures with herbicides from a different group when such use is permitted. Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices. Monitor treated weed populations for resistance development. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed. Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes. For further information or to report suspected resistance, contact Interprovincial Cooperative Ltd. at 1-204-233-3461.

GO TO SECTION 22

SECTION 20 - FIRST AID

1. **In case of poisoning**, call a physician or poison control centre IMMEDIATELY.
2. **If on skin or clothing**, take off contaminated clothing. Rinse skin IMMEDIATELY with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.
3. **If in eyes**, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.
4. **If swallowed**, call a poison control centre or doctor IMMEDIATELY for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.
5. **If inhaled**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.
6. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

GO TO SECTION 22

SECTION 21 - TOXICOLOGICAL INFORMATION

Treatment of any systemic intoxication should be primarily symptomatic and supportive. This product may cause severe irritation to the eyes. Overexposure to 2,4-D may cause coughing, burning, dizziness, or temporary loss of muscle coordination. Other possible effects of overexposure include fatigue, muscle weakness and nausea.

GO TO SECTION 22

SECTION 22 - RINSING AND DISPOSAL INFORMATION

For Recyclable Containers: Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site: 1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. 2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. **For Returnable Containers:** Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer). **For Returnable/Refillable Containers:** For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

(Base Label)

GROUP HERBICIDE

VITERRA

2,4-D Amine 600

LIQUID HERBICIDE

AGRICULTURAL

READ THE LABEL AND THE BOOKLET BEFORE USING.

GUARANTEE: 2,4-D (Present as Dimethylamine): 560 g a.e./L
Warning, contains the allergen soy.

REGISTRATION NO: 29248
PEST CONTROL PRODUCTS ACT

Net Contents: **10 litres**



Manufactured for:
Viterra Inc.
2625 Victoria Avenue
Regina, SK S4T 7T9
For Emergencies Call: 1-877-943-2876

By:
Interprovincial Cooperative Limited,
P.O. Box 1050, Saskatoon, Saskatchewan S7K 3M9
For Technical Support Call: 1-204-233-3461

VITERRA is a trademark of Viterra Inc.

RC 000-1208

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.